

# Conserved charges of higher $D$ Kerr–AdS spacetimes

S Deser<sup>1</sup>, Inanc Kanik<sup>2</sup> and Bayram Tekin<sup>2</sup>

<sup>1</sup> Department of Physics, Brandeis University, Waltham, MA 02454, USA

<sup>2</sup> Physics Department, Middle East Technical University, 06531 Ankara, Turkey

E-mail: [deser@brandeis.edu](mailto:deser@brandeis.edu), [ikanik@newton.physics.metu.edu.tr](mailto:ikanik@newton.physics.metu.edu.tr) and [btekin@metu.edu.tr](mailto:btekin@metu.edu.tr)

Received 9 June 2005, in final form 10 June 2005

Published 5 August 2005

Online at [stacks.iop.org/CQG/22/3383](http://stacks.iop.org/CQG/22/3383)

## Abstract

We compute the energy and angular momenta of recent  $D$ -dimensional Kerr–AdS solutions to cosmological Einstein gravity, using our invariant charge definitions.

PACS numbers: 04.20.–q, 04.20.Cv, 04.20.Ha, 11.30.–j

## 1. Introduction

Rotating solutions of cosmological Einstein gravity in  $D$  dimensions,  $R_{\mu\nu} = (D - 1)\Lambda g_{\mu\nu}$ , have been constructed recently [1, 2], extending earlier  $\Lambda = 0$  solutions of [3], themselves generalizations of the well-known  $D = 4$  metrics of [4] and [5], and of [6] in  $D = 5$ . These geometries provide a useful application of our recent generalized ‘conserved charge’ definitions, which are also extensions—of the original ADM [7], and AD [8] charges—to cover wider classes of actions [9, 11]: we will compute the energy and angular momenta of these new solutions.

Gravity theories have been historically endowed with a variety of seemingly different charge definitions, with different degrees of applicability and coordinate invariance. This topic has also seen much very recent activity, for example [12]. A summary and comparison of some of them is given in [13] which also includes a computation of the charges for Kerr–AdS black holes, using thermodynamic arguments; see also [14, 16]. Our results will agree with those, but we emphasize that in a general context, certain coincidences between charge definitions are suspect: for example, the frequently invoked ‘Komar’ charges, are in general not applicable, being highly gauge dependent [17].

## 2. Mass and angular momenta of Kerr–AdS

Let us briefly recapitulate the formulations of [8, 9]. The field equations of any metric model coupled to a (necessarily covariantly conserved) matter source  $\tau_{\mu\nu}$  are

$$\frac{\delta I}{\delta g_{\mu\nu}} \equiv \Phi_{\mu\nu}(g, R, \nabla R, \dots) = \kappa \tau_{\mu\nu}, \quad (1)$$

where  $\Phi_{\mu\nu}$  is an identically conserved tensor that can depend on curvatures and their derivatives. Decompose the metric into the sum of a background ‘vacuum’,  $\bar{g}_{\mu\nu}$  (which solves (1) for  $\tau_{\mu\nu} = 0$ ), plus a deviation  $h_{\mu\nu}$ , not necessarily small, that vanishes sufficiently rapidly far from the matter source:  $g_{\mu\nu} = \bar{g}_{\mu\nu} + h_{\mu\nu}$ . The field equations can be divided into a part linear in  $h_{\mu\nu}$  plus a non-linear remainder, which (with  $\tau_{\mu\nu}$ ) constitutes the total source  $T_{\mu\nu}$ . If the background  $\bar{g}_{\mu\nu}$  admits Killing vectors  $\bar{\xi}_\mu$ , obeying  $\bar{\nabla}_\mu \bar{\xi}_\nu + \bar{\nabla}_\nu \bar{\xi}_\mu = 0$ , then, up to normalization factors (which we shall fix later), the conserved Killing charges are

$$Q^\mu(\bar{\xi}) = \int_{\mathcal{M}} d^{D-1}x \sqrt{-\bar{g}} T^{\mu\nu} \bar{\xi}_\nu = \int_{\Sigma} dS_i \mathcal{F}^{\mu i}. \quad (2)$$

Here  $\Sigma$  is a  $D - 2$ -dimensional spacelike asymptotic hypersurface of the space  $\mathcal{M}$  and  $\mathcal{F}^{\mu i}$  is an anti-symmetric tensor, whose explicit form is model dependent. For Einstein’s theory with a cosmological constant,

$$Q^\mu = \frac{1}{4\Omega_{D-2}G_D} \int_{\Sigma} dS_i \{ \bar{\xi}_v \bar{\nabla}^\mu h^{iv} - \bar{\xi}_v \bar{\nabla}^i h^{\mu v} + \bar{\xi}^\mu \bar{\nabla}^i h - \bar{\xi}^i \bar{\nabla}^\mu h + h^{\mu\nu} \bar{\nabla}^i \bar{\xi}_v - h^{iv} \bar{\nabla}^\mu \bar{\xi}_v + \bar{\xi}^i \bar{\nabla}_v h^{\mu v} - \bar{\xi}^\mu \bar{\nabla}_v h^{iv} + h \bar{\nabla}^\mu \bar{\xi}^i \}, \quad (3)$$

where  $i$  takes values in  $1, 2, \dots, D - 2$  and the charge is normalized as shown, by dividing with the  $D$ -dimensional Newton’s constant and the solid angle. These charges are background gauge invariant under the diffeomorphisms  $\delta_\zeta h_{\mu\nu} = \bar{\nabla}_\mu \zeta_\nu + \bar{\nabla}_\nu \zeta_\mu$ :  $\delta_\zeta Q^\mu = 0$ .

Let us now calculate the conserved charges of the metrics [1] for  $D > 3$ . (We shall treat the special  $D = 3$  case at the end). They have the Kerr–Schild form [18, 19]

$$ds^2 = d\bar{s}^2 + \frac{2M}{U} (k_\mu dx^\mu)^2, \quad (4)$$

in terms of the de Sitter metric

$$d\bar{s}^2 = -W(1 - \Lambda r^2) dr^2 + F dr^2 + \sum_{i=1}^{N+\epsilon} \frac{r^2 + a_i^2}{1 + \Lambda a_i^2} d\mu_i^2 + \sum_{i=1}^N \frac{r^2 + a_i^2}{1 + \Lambda a_i^2} \mu_i^2 d\phi_i^2 + \frac{\Lambda}{W(1 - \Lambda r^2)} \left( \sum_{i=1}^{N+\epsilon} \frac{(r^2 + a_i^2) \mu_i d\mu_i}{1 + \Lambda a_i^2} \right)^2. \quad (5)$$

Here  $\epsilon = 0/1$  for odd/even dimensions and  $D = 2N + 1 + \epsilon$ . The null 1-form reads

$$k_\mu dx^\mu = F dr + W dr - \sum_{i=1}^N \frac{a_i \mu_i^2}{1 + \Lambda a_i^2} d\phi_i, \quad (6)$$

with

$$U \equiv r^\epsilon \sum_{i=1}^{N+\epsilon} \frac{\mu_i^2}{r^2 + a_i^2} \prod_{j=1}^N (r^2 + a_j^2), \quad W \equiv \sum_{i=1}^{N+\epsilon} \frac{\mu_i^2}{1 + \Lambda a_i^2}, \quad F \equiv \frac{1}{1 - \Lambda r^2} \sum_{i=1}^{N+\epsilon} \frac{r^2 \mu_i^2}{r^2 + a_i^2}. \quad (7)$$

To find the energy and angular momenta corresponding to (4), we must compute the charges  $Q^0$  for the corresponding Killing vectors: for the energy we shall take  $\bar{\xi}^\mu = (-1, 0)$  and each angular momentum has the appropriate unit entry  $(0, \dots, 1_i, \dots, 0)$ . Then

$$Q^0 = \frac{1}{4\Omega_{D-2}G_D} \int_{\Sigma} dS_r \{ g_{00} \bar{\nabla}^0 h^{r0} + g_{00} \bar{\nabla}^r h^{00} + h^{0v} \bar{\nabla}^r \bar{\xi}_v - h^{rv} \bar{\nabla}^0 \bar{\xi}_v + \bar{\nabla}_v h^{rv} \}, \quad (8)$$

Using the energy Killing vector, we obtain<sup>3</sup>

$$E_D = \frac{1}{4\Omega_{D-2}G_D} \int_{\Sigma} dS_r \left\{ g_{00}g^{rr}\partial_r h^{00} + \frac{1}{2}h^{00}g^{rr}\partial_r g_{00} - \frac{m}{U}g^{00}\partial_r g_{00} + 2m\partial_r U^{-1} \right. \\ \left. + \frac{2m}{U}g^{rr}\partial_r g_{rr} - \frac{m}{U}g^{rr}k^i k^j \partial_r g_{ij} + \frac{m}{U}g^{ij}\partial_r g_{ij} \right\}. \quad (9)$$

To compute  $E_D$ , one needs the large  $r$  behaviour of the integrand  $I$  of (9); since

$$g_{00} \rightarrow W\Lambda r^2, \quad F \rightarrow \frac{-1}{\Lambda r^2}, \quad U \rightarrow r^{D-3}, \quad k^\phi \rightarrow \frac{a_\phi}{r^2}, \quad (10)$$

then

$$I = \frac{2m}{r^{D-2}}[(D-1)W - 1]. \quad (11)$$

For completeness, let us also note how the determinant is calculated,

$$\det g = -W(1 - \Lambda r^2)F \prod_{i=1}^N \frac{(r^2 + a_i^2)\mu_i^2}{1 + \Lambda a_i^2} \det M, \quad (12)$$

Here  $M$  is the matrix representing the coefficients of the form  $d\mu_i d\mu_j$  in the metric, which can be expressed as (no repeated index summation)

$$M_{ij} = A_i \delta_{ij} + B_i B_j + C_i C_j \quad (13)$$

where

$$A_i = \frac{(r^2 + a_i^2)}{1 + \Lambda a_i^2}, \quad B_i = \sqrt{\frac{(r^2 + a_{N+\epsilon}^2)\mu_i}{1 + \Lambda a_{N+\epsilon}^2\mu_i}}, \quad (14)$$

$$C_i = \sqrt{\frac{\Lambda}{W(1 - \Lambda r^2)}} \left( \frac{(r^2 + a_i^2)}{1 + \Lambda a_i^2} - \frac{(r^2 + a_{N+\epsilon}^2)}{1 + \Lambda a_{N+\epsilon}^2} \right) \mu_i.$$

Then we have

$$\det M = \prod_{i=1}^{N+\epsilon-1} A_i \sum_{i=1}^{N+\epsilon-1} \left\{ \frac{B_i^2}{A_i} + \frac{C_i^2}{A_i} + \sum_{j \neq i} \frac{B_i^2 C_j^2}{A_i A_j} - \sum_{j \neq i} \frac{B_i B_j C_j C_i}{A_i A_j} \right\}. \quad (15)$$

Inserting (14) in the above equation, one gets

$$\det M = \frac{1}{W\mu_{N+\epsilon}^2} \prod_{i=1}^N \frac{1}{1 + \Lambda a_i^2}. \quad (16)$$

Using equations (16), (12), (11) the energy of the  $D$ -dimensional rotating black hole becomes

$$E_D = \frac{m}{\Xi} \sum_{i=1}^{\frac{D-1-\epsilon}{2}} \left\{ \frac{1}{\Xi_i} - (1 - \epsilon) \left( \frac{1}{2} \right) \right\}. \quad (17)$$

where

$$\Xi \equiv \prod_{i=1}^{\frac{D-1-\epsilon}{2}} (1 + \Lambda a_i^2), \quad \Xi_i \equiv 1 + \Lambda a_i^2. \quad (18)$$

<sup>3</sup> We are assuming that the background spacetime is AdS rather than dS, whose cosmological horizon causes complications. Some of these issues were addressed in [8, 9]. For details of acceptable asymptotic falloff to (A)dS in various dimensions, we refer to [10].

This expression reduces to the standard limits  $a_i \rightarrow 0$  and  $\Lambda \rightarrow 0$ , and agrees (up to a constant factor) with those of [13–15].

The computation of angular momenta follows along similar lines. Consider a given, say that  $i$ th (which we call the  $\phi$ ) component, i.e., the Killing vector  $\xi_{(i)}^\mu = (0, \dots, 0, 1_i, 0, \dots)$ . Then the corresponding Killing charge becomes

$$\begin{aligned} Q^0 &= \frac{1}{4\Omega_{D-2}G_D} \int_{\Sigma} dS_r \{g_{\phi\phi} \hat{\nabla}^0 h^{r\phi} - g_{\phi\phi} \hat{\nabla}^r h^{0\phi} + h^{0v} \hat{\nabla}^r \xi_v - h^{rv} \hat{\nabla}^0 \xi_v\} \\ &= \frac{1}{4\Omega_{D-2}G_D} \int_{\Sigma} dS_r \{-g_{\phi\phi} g^{rr} g^{00} \partial_r h_0^\phi\}. \end{aligned} \quad (19)$$

Once again the integrand can be calculated to be

$$I = \frac{(D-1)2ma_i\mu_i^2}{r^{D-2}(1+\Lambda a_i^2)}. \quad (20)$$

Putting the pieces together, the angular momentum is

$$J_i = \frac{ma_i}{\Xi\Xi_i}. \quad (21)$$

This expression again agrees with the previous ones. Note that, unlike in the energy expression,  $\epsilon$  does not appear here since even-dimensional spaces have as many independent 2-planes as the odd-dimensional spaces with one lower dimension<sup>4</sup>.

Having computed the desired conserved charges (17), (21) for Kerr–AdS spacetimes in  $D > 3$ , let us briefly turn our attention to the  $D = 3$  BTZ black hole [20]. This solution has long been studied but we recompute the charges with our method for the sake of completeness. The BTZ black hole differs from its higher dimensional counterparts in one very important aspect: for it, AdS is not the correct vacuum background [20]. The full metric is

$$ds^2 = (M - \Delta r^2) dt^2 + \frac{dr^2}{-M + \Delta r^2 + \frac{a^2}{4r^2}} - a dt d\phi + r^2 d\phi^2 \quad (22)$$

The background metric corresponds to  $M = 0$  and AdS corresponds to  $M = -1$ . Only AdS with  $J = 0$  is allowed for  $M < 0$ : the others have naked singularities. So we consider  $M > 0$  and compute the charges following our calculations above (about the  $M = 0$  background). We get the usual answers

$$E = M, \quad J = \Lambda. \quad (23)$$

BTZ black holes also solve the more general topologically massive gravity equations, where the Einstein term is augmented by the Cotton tensor [21],

$$G_{\mu\nu} + \Lambda g_{\mu\nu} + \frac{1}{\mu} C_{\mu\nu} = \kappa \tau_{\mu\nu}. \quad (24)$$

Conserved charges in this model were obtained in [22], in terms of those of the Einstein model  $Q_E^\mu$ ,

$$Q^\mu(\tilde{\xi}) = Q_E^\mu(\tilde{\xi}) + \frac{1}{\mu} \oint dS_i \{ \epsilon^{\mu\alpha\beta} \mathcal{G}_{\alpha\beta}^L \tilde{\xi}^\mu + \epsilon^{\nu\beta} \mathcal{G}_{\alpha\beta}^{\mu\beta} \tilde{\xi}_\nu + \epsilon^{\mu\nu\beta} \mathcal{G}_{\alpha\beta}^{Li} \tilde{\xi}_\nu \} + \frac{1}{\mu} Q_E^\mu(\epsilon \hat{\nabla} \tilde{\xi}), \quad (25)$$

where  $Q_E^\mu(\epsilon \hat{\nabla} \tilde{\xi})$  is the Einstein form but  $\tilde{\xi}$  is replaced with its curl. Once the contributions of the Cotton parts are computed the mass and the angular momentum of the BTZ black hole reads

$$E = M - \frac{\Delta a}{\mu}, \quad J = a - \frac{M}{\mu}, \quad (26)$$

a shift in values that may be compared with those of gravitational anyons [23], linearized solution of TMG but not of pure  $D = 3$  Einstein.

<sup>4</sup> For even dimensions, there is a nice relation between the energy and the angular momentum  $E = \sum_i \frac{J_i}{a_i}$ .

### 3. Mass and angular momenta in higher curvature models

We turn now to a slightly more formal exercise, which is to indicate the stability of our generic charge definition framework as it applies to a wider range of models, specifically higher derivative gravities. While Kerr-like solutions to  $R + R^2$  gravity models have yet to be discovered, it is not unlikely that they would approach the Einstein ones asymptotically. In that case, we could compute their conserved charges, defined as integrals at infinity, using the definitions for generic quadratic models [9]. Let us stick to the quadratic models of the form<sup>5</sup>

$$I = \int d^D x \sqrt{-g} \left\{ \frac{R}{2\kappa} + 2\Lambda_0 + \alpha R^2 + \beta R_{\mu\nu}^2 + \gamma (R_{\mu\nu\rho\sigma}^2 - 4R_{\mu\nu}^2 + R^2) \right\}. \quad (27)$$

This model allows constant curvature spacetimes with an effective cosmological constant given as

$$\Lambda = -\frac{1}{4f(\alpha, \beta, \gamma)\kappa} \{1 \pm \sqrt{1 + 8\kappa f(\alpha, \beta, \gamma)\Lambda_0}\} \quad \text{for } f(\alpha, \beta, \gamma) \neq 0, \quad (28)$$

where

$$f(\alpha, \beta, \gamma) = \frac{(D-4)}{(D-2)^2} (D\alpha + \beta) + \frac{\gamma(D-4)(D-3)}{(D-2)(D-1)}. \quad (29)$$

When the bare cosmological constant vanishes ( $\Lambda_0 = 0$ ), (A)dS spaces are still allowed and one has the + sign branch in (28). Conserved charges in this model, which we quote below, were defined in [9]

$$\begin{aligned} Q^\mu(\bar{\xi}) = & \left\{ \frac{1}{\kappa} + \frac{4\Lambda D\alpha}{D-2} + \frac{4\Lambda\beta}{D-1} + \frac{4\Lambda\gamma(D-4)(D-3)}{(D-2)(D-1)} \right\} \int d^{D-1}x \sqrt{-\bar{g}} \bar{\xi}_\nu \mathcal{G}_L^{\mu\nu} \\ & + (2\alpha + \beta) \int dS_i \sqrt{-g} \{ \bar{\xi}^\mu \bar{\nabla}^i R_L + R_L \bar{\nabla}^\mu \bar{\xi}^i - \bar{\xi}^i \bar{\nabla}^\mu R_L \} \\ & + \beta \int dS_i \sqrt{-g} \{ \bar{\xi}_\nu \bar{\nabla}^i \mathcal{G}_L^{\mu\nu} - \bar{\xi}_\nu \bar{\nabla}^\mu \mathcal{G}_L^{i\nu} - \mathcal{G}_L^{\mu\nu} \bar{\nabla}^i \bar{\xi}_\nu + \mathcal{G}_L^{i\nu} \bar{\nabla}^\mu \bar{\xi}_\nu \}. \end{aligned} \quad (30)$$

where  $\mathcal{G}_L^{\mu\nu}$  and  $R_L$  are the linear parts of the Einstein tensor and the scalar curvature, respectively. The second and the third line vanish for Einstein spaces. The first line, on the other hand, is just a factor times the cosmological Einstein theory's charges (3). Therefore for asymptotic Kerr–AdS solutions, their conserved charges are given by the first term in (30), under the condition (28). Let us specifically consider the popular Einstein–Gauss–Bonnet theory,  $\alpha = \beta = 0$ . Also, implementing the condition (28) (with the + sign) we have

$$Q^\mu = -\sqrt{1 + 8\kappa f(\gamma, 0, 0)\Lambda_0} \frac{1}{\kappa} \int d^{D-1}x \sqrt{-\bar{g}} \bar{\xi}_\nu \mathcal{G}_L^{\mu\nu}. \quad (31)$$

Although the energy seems to have the wrong sign, this is a red herring: as shown in [24], for the non-rotating case the exact metric reads

$$ds^2 = g_{00} dt^2 + g_{rr} dr^2 + r^2 d\Omega_{D-2} \quad (32)$$

$$-g_{00} = g_{rr}^{-1} = 1 + \frac{r^2}{4\kappa\gamma(D-3)(D-4)} \left\{ 1 \pm \left[ 1 + 32\gamma\kappa(D-3)(D-4) \frac{m}{(D-2)r^{D-1}} \right]^{\frac{1}{2}} \right\}, \quad (33)$$

whose asymptotic forms branch into Schwarzschild and Schwarzschild–de-Sitter respectively,

$$-g_{00} = 1 - \frac{4m}{(D-2)r^{D-3}}, \quad -g_{00} = 1 + \frac{4m}{(D-2)r^{D-3}} + \frac{r^2}{\gamma(D-3)(D-4)}. \quad (34)$$

<sup>5</sup> Note that we changed normalization of the cosmological constant compared to the previous section.

We see that the SdS branch comes with the ‘wrong’ sign compared to the usual Schwarzschild one. Therefore, the minus sign in the energy becomes positive once  $\mathcal{G}_L^{\mu\nu}$  is explicitly computed. We conclude that the conserved charges in the Einstein–Gauss–Bonnet theory for such asymptotic solutions would be simply proportional to those of (17), (21) cosmological gravity [25]:

$$E_{\text{GB}} = \sqrt{1 + 8\kappa f(\gamma, 0, 0)\Lambda_0} E_D, \quad J_i(\text{GB}) = \sqrt{1 + 8\kappa f(\gamma, 0, 0)\Lambda_0} J_i, \quad (35)$$

It is important to note that if the coefficient  $\sqrt{1 + 8\kappa f(\gamma, 0, 0)\Lambda_0}$  does not vanish, then one can simply rescale<sup>6</sup> the Killing charges to get the Einstein charges (17), (21).

#### 4. Conclusions

Using the charge definitions via background Killing charges of [8, 9] we have computed the mass and angular momenta of the rotating Kerr–AdS black holes for  $D$  dimensions for cosmological Einstein gravity. As a test of stability, we checked that the corresponding charge definitions for higher order would lead to the same values for asymptotically similar geometries up to the indicated constant rescaling.

#### Acknowledgments

The work of SD is supported by NSF grant no PHY 04-01667; that of BT by the ‘Young Investigator Fellowship’ of Turkish Academy of Sciences (TUBA) and by a TUBITAK Kariyer grant. BT thanks Özgür Sarioğlu for a fruitful discussion on TMG.

#### References

- [1] Gibbons G W, Lu H, Page D N and Pope C N 2005 The general kerr–de sitter metrics in all dimensions *J. Geom. Phys.* **53** 49
- [2] Gibbons G W, Lu H, Page D N and Pope C N 2004 Rotating black holes in higher dimensions with a cosmological constant *Phys. Rev. Lett.* **93** 171102
- [3] Myers R C and Perry M J 1986 Black holes in higher dimensional space-times *Ann. Phys.* **172** 304
- [4] Carter B 1968 Hamilton–Jacobi and Schrödinger separable solutions of Einstein’s equations *Commun. Math. Phys.* **10** 280
- [5] Kerr R P 1963 Gravitational field of a spinning mass as an example of algebraically special metrics *Phys. Rev. Lett.* **11** 237
- [6] Hawking S W, Hunter C J and Taylor-Robinson M M 1999 Rotation and the adS/CFT correspondence *Phys. Rev. D* **59** 064005
- [7] Arnowitt R, Deser S and Misner C 1959 *Phys. Rev.* **116** 1322  
Arnowitt R, Deser S and Misner C 1960 *Phys. Rev.* **117** 1595  
Arnowitt R, Deser S and Misner C 1962 *Gravitation: An Introduction to Current Research* ed L Witten (New York: Wiley) (Preprint gr-qc/0405109)
- [8] Abbott L F and Deser S 1982 Stability of gravity with a cosmological constant *Nucl. Phys. B* **195** 76
- [9] Deser S and Tekin B 2003 Energy in generic higher curvature gravity theories *Phys. Rev. D* **67** 084009
- [10] Henneaux M and Teitelboim C 1985 Asymptotically anti-de Sitter spaces *Commun. Math. Phys.* **98** 391  
Henneaux M 1985 Asymptotically anti-de Sitter universes in  $D = 3, 4$  and higher dimensions *Proc. General Relativity (Rome 1985)* Part B pp 959–66  
Brown J D and Henneaux M 1986 Central charges in the canonical realization of asymptotic symmetries: an example from three-dimensional gravity *Commun. Math. Phys.* **104** 207
- [11] Deser S and Tekin B 2002 Gravitational energy in quadratic curvature gravities *Phys. Rev. Lett.* **89** 101101
- [12] Petrov A N 2005 A note on the Deser–Tekin charges *Class. Quantum Grav.* **22** L83 (Preprint gr-qc/0504058)

<sup>6</sup> To avoid rescaling of the Killing charges, one could appropriately choose the perturbation branch. We thank A N Petrov for his remarks on this.

- [13] Gibbons G W, Perry M J and Pope C N 2005 The first law of thermodynamics for Kerr–anti-de Sitter black holes *Class. Quantum Grav.* **22** 1503 (*Preprint* hep-th/0408217)
- [14] Deruelle N and Katz J 2005 On the mass of a Kerr–anti-de Sitter spacetime in  $D$  dimensions *Class. Quantum Grav.* **22** 421
- [15] Barnich G and Compere G 2005 Generalized Smarr relation for Kerr AdS black holes from improved surface integrals *Phys. Rev. D* **71** 044016
- [16] Katz J, Bicak J and Lynden-Bell D 1997 Relativistic conservation laws and integral constraints for large cosmological perturbations *Phys. Rev. D* **55** 5957
- [17] Misner C W 1963 Gravitational field energy and  $g_{00}$  *Phys. Rev.* **130** 1590
- [18] Kerr R P and Schild A 1965 Some algebraically degenerate solutions of Einstein’s gravitational field equations *Proc. Symp. Appl. Math.* **17** 199
- [19] Gurses M and Gursey F 1975 Lorentz invariant treatment of Kerr–Schild geometry *J. Math. Phys.* **16** 2385
- [20] Banados M, Teitelboim C and Zanelli J 1992 The black hole in three-dimensional space-time *Phys. Rev. Lett.* **69** 1849
- [21] Deser S, Jackiw R and Templeton S 1982 Topologically massive gauge theories *Ann. Phys.* **140** 372  
Deser S, Jackiw R and Templeton S 1982 Three-dimensional massive gauge theories *Phys. Rev. Lett.* **48** 975
- [22] Deser S and Tekin B 2003 Energy in topologically massive gravity *Class. Quantum Grav.* **20** L259
- [23] Deser S 1990 Gravitational anyons *Phys. Rev. Lett.* **64** 611
- [24] Boulware D G and Deser S 1985 String generated gravity models *Phys. Rev. Lett.* **55** 2656
- [25] Deruelle N and Morisawa Y 2005 Mass and angular momenta of Kerr anti-de Sitter spacetimes in Einstein–Gauss–Bonnet theory *Class. Quantum Grav.* **22** 933